
## Number and Proportion of AIDS Cases among Female Adults and Adolescents 1985–2004—United States



Year of diagnosis






# AIDS Cases and Rates for Female Adults and Adolescents, by Race/Ethnicity 2004—50 States and D.C. Rate (per

| Race/Ethnicity                    | <u>Number</u> | %  | 100,000<br>population) |
|-----------------------------------|---------------|----|------------------------|
| White, not Hispanic               | 1,860         | 17 | 2.1                    |
| Black, not Hispanic               | 7,395         | 67 | 48.2                   |
| Hispanic                          | 1,643         | 15 | 11.1                   |
| Asian/Pacific Islander            | 92            | <1 | 1.6                    |
| American Indian/<br>Alaska Native | 64            | <1 | 6.4                    |
| Total*                            | 11,109        |    | 9.0                    |






### Proportion of AIDS Cases among Female Adults and Adolescents, by Transmission Category 2004—United States







<sup>\*\*</sup>Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.



### Proportion of AIDS Cases among Female Adults and Adolescents, by Transmission Category and Age at Diagnosis, 2004—United States

Age at diagnosis (in years)

| Transmission category           | 13–19         | 20–24 | 25 <b>–</b> 34 | 35 <b>–</b> 44 | <b>≥</b> 45 |
|---------------------------------|---------------|-------|----------------|----------------|-------------|
|                                 | <del></del> % | %     | <del></del> %  |                | %           |
| Injection drug use              | 12            | 18    | 21             | 31             | 31          |
| High risk heterosexual contact* | 66            | 79    | 77             | 67             | 67          |
| Other/not identified**          | 23            | 3     | 2              | 2              | 2           |

Note. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.


<sup>\*\*</sup>Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.





<sup>\*</sup>Heterosexual contact with a person known to have or at high risk for HIV infection.


### AIDS Rates for Female Adults and Adolescents, Reported in 2004—United States



<sup>\*</sup> Rates were not calculated for states that reported fewer than 5 AIDS cases in females in 2004.




### AIDS Cases among Female Adults and Adolescents Attributed to Injection Drug Use or High Risk Heterosexual Contact, by Region, 2000–2004—50 States and D.C.








### Reported AIDS Cases among Female Adults and Adolescents, by Region and Race/Ethnicity, 2004 50 States and D.C.







#### Female Adults and Adolescents 15 to 44 Years of Age Reported to be Living with HIV Infection (not AIDS), 2004—42 Areas






Note. Data from 42 areas with confidential name-based HIV infection reporting as of December 2004. Data based on person's age as of December 31, 2004. \* Includes 303 persons reported from areas with confidential name-based HIV infection reporting, but who were residents of areas without confidential name-based HIV infection reporting. Includes 173 persons whose state of residence is unknown or missing.

\*\* Pennsylvania implemented reporting only for areas outside the city of Philadelphia.




Female Adults and Adolescents 15 to 44 Years of Age Reported to be Living with AIDS, 2004—United States







### Proportion of HIV/AIDS Cases among Female Adults and Adolescents, by Transmission Category 2004—33 States



Note. Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2000. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.






<sup>\*</sup>Heterosexual contact with a person known to have or at high risk for HIV infection.

<sup>\*\*</sup>Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

### Proportion of HIV/AIDS Cases and Population among Female Adults and Adolescents, by Race/Ethnicity 2004—33 States







## Proportion of HIV/AIDS Cases among Female Adults and Adolescents, by Transmission Category and Age at Diagnosis, 2004—33 States

|                                 |       | Age at d | iagnosis          | (in years) |     |
|---------------------------------|-------|----------|-------------------|------------|-----|
| Transmission category           | 13-19 | 20-24    | <u>25-34</u><br>% | 35-44      | 45+ |
| Injection drug use              | 15    | 15       | 18                | 24         | 23  |
| High risk heterosexual contact* | 83    | 84       | 81                | 75         | 75  |
| Other/not identified**          | 1     | 1        | 1                 | 2          | 2   |


Note. Data include persons with a diagnosis of HIV regardless of AIDS status at diagnosis. Data from 33 states with confidential named-based HIV infection reporting since at least 2000. Data have been adjusted for reporting delays. \* Heterosexual contact with a person known to have or at high risk for HIV infection.

<sup>\*\*</sup> Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.





### Diagnosis Rates of HIV/AIDS for Female Adults and Adolescents, 2004—33 States







### Estimated Number of HIV/AIDS Cases and Rates for Female Adults and Adolescents, by Race/Ethnicity 2004—33 States

Pata

| Race/Ethnicity                    | Cases* | (cases per 100,000 population) |
|-----------------------------------|--------|--------------------------------|
| White, not Hispanic               | 1,782  | 3.2                            |
| Black, not Hispanic               | 7,009  | 67.0                           |
| Hispanic                          | 1,400  | 16.3                           |
| Asian/Pacific Islander            | 94     | 4.1                            |
| American Indian/<br>Alaska Native | 57     | 7.7                            |
| Total**                           | 10,391 | 13.2                           |

Note. Data include persons with a diagnosis of HIV regardless of AIDS status at diagnosis. Data from 33 states with confidential named-based HIV infection reporting since at least 2000. Data have been adjusted for reporting delays. \* Data exclude persons from US dependencies, possessions and associated nations because of the lack of data by race from these areas.

<sup>\*\*</sup> Includes 49 female adults and adolescents of unknown race or multiple races.

